arXiv:1012.2681v2 [math.NT] 14 Oct 2012

WZ-PROOFS OF “DIVERGENT” RAMANUJAN-TYPE SERIES

JESÚS GUILLERA

Dedicated to Herb Wilf on his 80^{th} birthday

Abstract. We prove some “divergent” Ramanujan-type series for $1/\pi$ and $1/\pi^2$ applying a Barnes-integrals strategy of the WZ-method.

1. Wilf-Zeilberger’s pairs

We recall that a function $A(n,k)$ is *hypergeometric* in its two variables if the quotients

$$\frac{A(n+1,k)}{A(n,k)} \quad \text{and} \quad \frac{A(n,k+1)}{A(n,k)}$$

are rational functions in n and k, respectively. Also, a pair of hypergeometric functions in its two variables, $F(n,k)$ and $G(n,k)$, is said to be a *Wilf and Zeilberger (WZ) pair* [13, Chapt. 7] if

$$F(n+1,k)-F(n,k)=G(n,k+1)-G(n,k). \tag{1}$$

In this case, H. S. Wilf and D. Zeilberger [17] have proved that there exists a rational function $C(n,k)$ such that

$$G(n,k)=C(n,k)F(n,k). \tag{2}$$

The rational function $C(n,k)$ is the so-called *certificate* of the pair (F,G). To discover WZ-pairs, we use Zeilberger’s Maple package EKHAD [13, Appendix A]. If EKHAD certifies a function, we have found a WZ-pair! We will write the functions $F(n,k)$ and $G(n,k)$ using rising factorials, also called Pochhammer symbols, rather than the ordinary factorials. The rising factorial is defined by

$$(x)_n = \begin{cases} x(x+1)\cdots(x+n-1), & n \in \mathbb{Z}^+, \\ 1, & n=0, \end{cases} \tag{3}$$

or more generally by $(x)_t=\Gamma(x+t)/\Gamma(x)$. For $t \in \mathbb{Z}-\mathbb{Z}^-$, this last definition coincide with (3). But it is more general because it is also defined for all complex x and t such that $x+t \in \mathbb{C}-(\mathbb{Z}-\mathbb{Z}^+)$.

Key words and phrases. Hypergeometric series; WZ-method; Ramanujan-type series for $1/\pi$ and $1/\pi^2$; Barnes Integrals.

2. A Barnes-integrals WZ strategy

If we sum (1) over all $n \geq 0$, we get

$$\sum_{n=0}^{\infty} G(n,k) - \sum_{n=0}^{\infty} G(n,k+1) = -F(0,k) + \lim_{n \to \infty} F(n,k) \tag{4}$$

whenever the series above are convergent and the limit is finite. D. Zeilberger was the first to apply the WZ-method to prove a Ramanujan-type series for $1/\pi$ [4]. Following his idea, in a series of papers [5], [6], [9], [10] and in the author's thesis [8], we use WZ-pairs together with formula (4) to prove a total of eleven Ramanujan-type series for $1/\pi$ and four Ramanujan-like series for $1/\pi^2$. However, while we discovered those pairs we also found some WZ-pairs corresponding to "divergent" Ramanujan-type series [12], like the following pair:

$$F(n,k) = A(n,k) \frac{(-1)^n}{\Gamma(n+1)} \left(\frac{16}{9} \right)^n, \quad G(n,k) = B(n,k) \frac{(-1)^n}{\Gamma(n+1)} \left(\frac{16}{9} \right)^n,$$

where

$$A(n,k) = U(n,k) \frac{-n(n-2)}{3(n+2k+1)}, \quad B(n,k) = U(n,k)(5n+6k+1),$$

and

$$U(n,k) = \frac{\left(\frac{1}{2}\right)_n \left(\frac{1}{4} + \frac{3k}{2}\right)_n \left(\frac{3}{4} + \frac{3k}{2}\right)_n}{(1+k)_n (1+2k)_n} \frac{\left(\frac{1}{6}\right)_k \left(\frac{5}{6}\right)_k}{(1)_k^2}.$$

We cannot use formula (4) with this pair because the series is divergent and the limit is infinite, due to the factor $(-16/9)^n$. To deal with this kind of WZ-pairs we will proceed as follows: First we replace the factor $(-1)^n$ with $\Gamma(n+1)\Gamma(-n)$. By doing it we again get a WZ-pair, because $(-1)^n$ and $\Gamma(n+1)\Gamma(-n)$ transform formally in the same way under the substitution $n \to n+1$; namely, the sign changes. To fix ideas, the modified version of the WZ-pair above is

$$\widetilde{F}(s,t) = A(s,t)\Gamma(-s) \left(\frac{16}{9} \right)^s, \quad \widetilde{G}(s,t) = B(s,t)\Gamma(-s) \left(\frac{16}{9} \right)^s.$$

Then, integrating from $s = -i\infty$ to $s = i\infty$ along a path $\mathcal{P}$ (curved if necessary) which separates the poles of the form $s = 0, 1, 2 \ldots$ from all the other poles, we obtain

$$\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} B(s,t)\Gamma(-s)(-z)^s ds = \sum_{n=0}^{\infty} B(n,t) \frac{z^n}{n!}, \quad |z| < 1, \tag{5}$$

where we have used the Barnes integral theorem, which is an application of Cauchy's residues theorem using a contour which closes the path with a right side semicircle of center at the origin and infinite radius. The Barnes integral gives the analytic continuation of the series to $z \in \mathbb{C} - [1, \infty)$. Integrating along the same path the identity $\widetilde{G}(s,t+1)-\widetilde{G}(s,t)=\widetilde{F}(s+1,t)-\widetilde{F}(s,t)$, we obtain

$$\begin{aligned}(6)\qquad \int_{-i\infty}^{i\infty}\widetilde{G}(s,t+1)ds-\int_{-i\infty}^{i\infty}\widetilde{G}(s,t)ds&=\int_{-i\infty}^{i\infty}\widetilde{F}(s+1,t)ds-\int_{-i\infty}^{i\infty}\widetilde{F}(s,t)ds\\ =\int_{1-i\infty}^{1+i\infty}\widetilde{F}(s,t)ds-\int_{-i\infty}^{i\infty}\widetilde{F}(s,t)ds&=-\int_{\mathcal{C}}\widetilde{F}(s,t)ds,\end{aligned}$$

where $\mathcal{C}$ is the contour limited by the path $\mathcal{P}$, the same path but moved one unit to the right, and the lines $y=-\infty$ and $y=+\infty$. As the only pole inside this contour is at $s=0$ and the residue at this point is zero, the last integral is zero and we have

$$(7)\qquad \int_{-i\infty}^{i\infty}\widetilde{G}(s,t)ds=\int_{-i\infty}^{i\infty}\widetilde{G}(s,t+1)ds.$$

This implies, by Weierstrass's theorem [16], that

$$\begin{aligned}\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\widetilde{G}(s,t)ds&=\lim_{t\to\infty}\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\widetilde{G}(s,t)ds=\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\lim_{t\to\infty}\widetilde{G}(s,t)ds\\ &=\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\frac{3}{\pi}\left(\frac{1}{2}\right)_s\Gamma(-s)2^s ds=\frac{\sqrt{3}}{\pi},\end{aligned}$$

where the last equality holds because

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\left(\frac{1}{2}\right)_s\Gamma(-s)(-z)^s ds=\sum_{n=0}^{\infty}\frac{\left(\frac{1}{2}\right)_n}{(1)_n}z^n=\frac{1}{\sqrt{1-z}},\qquad |z|<1,$$

implies that

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\left(\frac{1}{2}\right)_s\Gamma(-s)(-z)^s ds=\frac{1}{\sqrt{1-z}},\qquad z\in\mathbb{C}-[1,\infty).$$

Hence, we have

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\frac{\left(\frac{1}{2}\right)_s\left(\frac{1}{4}+\frac{3t}{2}\right)_s\left(\frac{3}{4}+\frac{3t}{2}\right)_s}{(1+t)_s(1+2t)_s}\frac{\left(\frac{1}{6}\right)_t\left(\frac{5}{6}\right)_t}{(1)_t^2}(5s+6t+1)\Gamma(-s)\left(\frac{4}{3}\right)^{2s}ds=\frac{\sqrt{3}}{\pi},$$

or equivalently

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\frac{\left(\frac{1}{2}\right)_s\left(\frac{1}{4}+\frac{3t}{2}\right)_s\left(\frac{3}{4}+\frac{3t}{2}\right)_s}{(1+t)_s(1+2t)_s}(5s+6t+1)\Gamma(-s)\left(\frac{4}{3}\right)^{2s}ds=\frac{\sqrt{3}}{\pi}\frac{(1)_t^2}{\left(\frac{1}{6}\right)_t\left(\frac{5}{6}\right)_t}.$$

Finally, substituting $t=0$, we see that

$$(8)\qquad \frac{1}{2\pi i}\int_{-i\infty}^{i\infty}\frac{\left(\frac{1}{2}\right)_s\left(\frac{1}{4}\right)_s\left(\frac{3}{4}\right)_s}{(1)_s^2}(5s+1)\Gamma(-s)\left(\frac{4}{3}\right)^{2s}ds=\frac{\sqrt{3}}{\pi}.$$

It is very convenient to write the Barnes integral in hypergeometric notation. By the definition of hypergeometric series, we see that for $-1\le z<1$, we have

$$\sum_{n=0}^{\infty}\frac{\left(\frac{1}{2}\right)_n(s)_n(1-s)_n}{(1)_n^3}z^n={}_3F_2\left(\begin{matrix}\frac{1}{2}, & s, & 1-s\\ & 1, & 1\end{matrix}\,\middle|\,z\right)$$

and

$$\sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n (s)_n (1-s)_n}{(1)_n^3} n z^n = \frac{1}{2} s(1-s) z \, {}_3F_2\left(\begin{matrix} \frac{3}{2}, & 1+s, & 2-s \\ & 2, & 2 \end{matrix} \,\middle|\, z \right),$$

where the notation on the right side stands for the analytic continuation of the series on the left. Hence, we can write (8) in the form

$${}_3F_2\left(\begin{matrix} \frac{1}{2}, & \frac{1}{4}, & \frac{3}{4} \\ & 1, & 1 \end{matrix} \,\middle|\, \frac{-16}{9} \right) - \frac{5}{6} \, {}_3F_2\left(\begin{matrix} \frac{3}{2}, & \frac{5}{4}, & \frac{7}{4} \\ & 2, & 2 \end{matrix} \,\middle|\, \frac{-16}{9} \right) = \frac{\sqrt{3}}{\pi}.$$

If, instead of integrating to the right side, we integrate (8) along a contour which closes the path $\mathcal{P}$ with a semicircle of center $s=0$ taken to the left side with an infinite radius, then we have poles at $s=-n-1/2$, at $s=-n-1/4$ and at $s=-n-3/4$ for $n=0,1,2,\dots$, and we obtain

$$\begin{aligned} &\frac{\sqrt{3}}{2} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{2}\right)_n^3}{(1)_n \left(\frac{3}{4}\right)_n \left(\frac{5}{4}\right)_n} (10n+3)(-1)^n \left(\frac{3}{4}\right)^{2n} \\ &\qquad - \frac{\sqrt{2}\,\pi^2}{8\,\Gamma\left(\frac{3}{4}\right)^4} \sum_{n=0}^{\infty} \frac{\left(\frac{1}{4}\right)_n^3}{(1)_n \left(\frac{1}{2}\right)_n \left(\frac{3}{4}\right)_n} (20n+1)(-1)^n \left(\frac{3}{4}\right)^{2n} \\ &\qquad\quad - \frac{3\sqrt{2}\,\Gamma\left(\frac{3}{4}\right)^4}{16\,\pi^2} \sum_{n=0}^{\infty} \frac{\left(\frac{3}{4}\right)_n^3}{(1)_n \left(\frac{3}{2}\right)_n \left(\frac{5}{4}\right)_n} (20n+11)(-1)^n \left(\frac{3}{4}\right)^{2n} = 1. \end{aligned}$$

which is an identity relating three convergent series.

3. Other examples

In a similar way we can prove other identities of the same kind, for example,

$$\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{\left(\frac{1}{2}+t\right)_s^3 \left(\frac{1}{2}\right)_s^2}{(1+t)_s^3 (1+2t)_s} (10s^2+6s+1+14st+4t^2+4t) \Gamma(-s) 2^{2s} ds = \frac{4}{\pi^2} \frac{(1)_t^4}{\left(\frac{1}{2}\right)_t^4},$$

$$\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{\left(\frac{1}{2}\right)_s \left(\frac{1}{2}+t\right)_s^2}{(1)_s (1+2t)_s} (3s+2t+1) \Gamma(-s) 2^{3s} ds = \frac{1}{\pi} \frac{(1)_t}{\left(\frac{1}{2}\right)_t},$$

and

$$\begin{aligned} &\frac{1}{2\pi i} \int_{-i\infty}^{i\infty} \frac{\left(\frac{1}{2}\right)_s \left(\frac{1}{2}+2t\right)_s \left(\frac{1}{3}+t\right)_s \left(\frac{2}{3}+t\right)_s}{\left(\frac{1}{2}+\frac{t}{2}\right)_s \left(1+\frac{t}{2}\right)_s (1+t)_s} \\ &\qquad \times \frac{(15s+4)(2s+1)+t(33s+16)}{2s+t+1} \Gamma(-s) 2^{2s} ds = \frac{3\sqrt{3}}{\pi} \frac{1}{2^{6t}} \frac{(1)_t^2}{\left(\frac{1}{4}\right)_t \left(\frac{3}{4}\right)_t}. \end{aligned}$$

In the two last examples the hypothesis of Weierstrass theorem fail and hence we cannot apply it, but we obtain the sum using Meurman's periodic version of Carlson's theorem [2, p. 39] which asserts that if $H(z)$ is a periodic entire function of period 1 and there is a real number $c < 2\pi$ such that $H(z) = \mathcal{O}(\exp(c|Im(z)|))$ for all $z \in \mathbb{C}$, then $H(z)$ is constant [1, Appendix] and [11, Thm. 2.3]. In the second and third examples we determine the constants $1/\pi$ and $3\sqrt{3}/\pi$ taking $t = 1/2$ and $t = -1/3$ respectively. Substituting $t = 0$ in the above examples, we obtain respectively

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty} \frac{\left(\frac{1}{2}\right)_s^5}{(1)_s^4}(10s^2+6s+1)\Gamma(-s)2^{2s}ds = \frac{4}{\pi^2}, \tag{9}$$

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty} \frac{\left(\frac{1}{2}\right)_s^3}{(1)_s^2}(3s+1)\Gamma(-s)2^{3s}ds = \frac{1}{\pi}, \tag{10}$$

and

$$\frac{1}{2\pi i}\int_{-i\infty}^{i\infty} \frac{\left(\frac{1}{2}\right)_s\left(\frac{1}{3}\right)_s\left(\frac{2}{3}\right)_s}{(1)_s^2}(15s+4)\Gamma(-s)2^{2s}ds = \frac{3\sqrt{3}}{\pi}. \tag{11}$$

Using hypergeometric notation, we can write (9), (10) and (11) respectively in the following forms:

$$\begin{aligned}{}_5F_4\left(\begin{matrix}\frac12, & \frac12, & \frac12, & \frac12, & \frac12\\ & 1, & 1, & 1, & 1\end{matrix}\,\middle|\,-4\right) - \frac{3}{4}\,{}_5F_4\left(\begin{matrix}\frac32, & \frac32, & \frac32 & \frac32 & \frac32\\ & 2, & 2, & 2, & 2\end{matrix}\,\middle|\,-4\right)&\\ -\frac{5}{4}\,{}_5F_4\left(\begin{matrix}\frac32, & \frac32, & \frac32 & \frac32 & \frac32\\ & 2, & 2, & 2, & 1\end{matrix}\,\middle|\,-4\right) &= \frac{4}{\pi^2},\end{aligned}$$

$${}_3F_2\left(\begin{matrix}\frac12, & \frac12, & \frac12\\ & 1, & 1\end{matrix}\,\middle|\,-8\right) - 3\,{}_3F_2\left(\begin{matrix}\frac32, & \frac32, & \frac32\\ & 2, & 2\end{matrix}\,\middle|\,-8\right) = \frac{1}{\pi},$$

and

$$4\,{}_3F_2\left(\begin{matrix}\frac12, & \frac13, & \frac23\\ & 1, & 1\end{matrix}\,\middle|\,-4\right) - \frac{20}{3}\,{}_3F_2\left(\begin{matrix}\frac32, & \frac43, & \frac53\\ & 2, & 2\end{matrix}\,\middle|\,-4\right) = \frac{3\sqrt{3}}{\pi}.$$

Related applications of the WZ-method for Barnes-type integrals are for example in [3, Sect. 5.2] and [14].

4. The dual of a “divergent” Ramanujan-type series

The WZ duality technique [13, Ch. 7] allows to transform pairs which lead to divergences into pairs which lead to convergent series. To get the dual $\widehat{G}(n,k)$ of $G(-n,-k)$, we make the following changes:

$$(a)_{-n} \to \frac{(-1)^n}{(1-a)_n}, \quad (1)_{-n} \to \frac{n(-1)^n}{(1)_n}, \quad (a)_{-k} \to \frac{(-1)^k}{(1-a)_k}, \quad (1)_{-k} \to \frac{k(-1)^k}{(1)_k}.$$

Example 1. The package EKHAD certifies the pair

$$F(n,k)=U(n,k)\frac{2n^2}{2n+k}, \qquad G(n,k)=U(n,k)\frac{6n^2+2n+k+4nk}{2n+k}, \tag{12}$$

where

$$U(n,k)=\frac{\left(\frac{1}{2}\right)_n^2\left(1+\frac{k}{2}\right)_n\left(\frac{1}{2}+\frac{k}{2}\right)_n}{(1)_n^2(1+k)_n^2}\frac{\left(\frac{1}{2}\right)_k}{(1)_k}4^n=\frac{(2n)!^2(2n+k)!(2k)!}{n!^4k!(n+k)!^2}\frac{1}{16^n4^k}.$$

We cannot use this WZ-pair to obtain a Ramanujan-like evaluation because, as $z>1$, the corresponding series and also the corresponding Barnes integral are both divergent. However, we will see how to use it to evaluate a related convergent series. What we will do is to apply the WZ duality technique. Thus, if we take the dual of $G(-n,-k)$ and replace k with $k-1$, we obtain

$$\widehat{G}(n,k)=\frac{1}{U(n,k)}\frac{2(2k-1)(2n+k)}{n^2(n+k)^2(n+k-1)^2}(6n^2-6n+1-k+4nk),$$

and EHKAD finds its companion

$$\widehat{F}(n,k)=\frac{1}{U(n,k)}\frac{-2(2n+k)(2n+k-1)(2n-1)^2}{n^2(n+k)^2(n+k-1)^2}.$$

Applying Zeilberger's formula

$$\sum_{n=j}^{\infty}(\widehat{F}(n+1,n)+\widehat{G}(n,n))=\sum_{n=j}^{\infty}\widehat{G}(n,j)$$

with $j=1$, we obtain

$$\sum_{n=1}^{\infty}\left(\frac{16}{27}\right)^n\frac{(1)_n^3}{\left(\frac{1}{2}\right)_n\left(\frac{1}{3}\right)_n\left(\frac{2}{3}\right)_n}\frac{11n-3}{n^3}=16\sum_{n=1}^{\infty}\frac{1}{4^n}\frac{(1)_n^3}{\left(\frac{1}{2}\right)_n^3}\frac{3n-1}{n^3}. \tag{13}$$

The series in (13) are dual to Ramanujan-type "divergent" series, and in [7, p. 221] we proved that the series on the right side is equal to $\pi^2/2$. Hence

$$\sum_{n=1}^{\infty}\left(\frac{16}{27}\right)^n\frac{(1)_n^3}{\left(\frac{1}{2}\right)_n\left(\frac{1}{3}\right)_n\left(\frac{2}{3}\right)_n}\frac{11n-3}{n^3}=8\pi^2. \tag{14}$$

Formula (14), as well as other similar formulas, was conjectured in [15, Conj 1.4] by Zhi-Wei Sun.

Example 2. The package EKHAD certifies the pair

$$F(n,k)=U(n,k)\frac{64n^3}{(2k+1)(2n-2k+1)},$$

$$G(n,k)=U(n,k)\frac{(2n+1)^2(11n+3)-12k(2n^2+3nk+n+k)}{(2n+1)^2},$$

where

$$U(n,k)=\frac{\left(\frac{1}{2}-k\right)_n\left(\frac{1}{2}+k\right)_n^2\left(\frac{1}{3}\right)_n\left(\frac{1}{3}\right)_n}{(1)_n^3\left(\frac{1}{2}\right)_n^2}\left(\frac{27}{16}\right)^n.$$

Taking the dual $\widehat{G}(n,k)$ of $G(-n,-k)$, replacing n with $n+x$ and applying Zeilberger's theorem

$$\sum_{n=0}^{\infty}\widehat{G}(n+x,0)=\lim_{k\to\infty}\sum_{n=0}^{\infty}\widehat{G}(n+x,k)+\sum_{k=0}^{\infty}\widehat{F}(x,k),$$

where $\widehat{F}(n,k)$ is the companion of $\widehat{G}(n,k)$, we obtain

$$\sum_{n=0}^{\infty}\frac{(1+x)_n^3}{\left(\frac{1}{2}+x\right)_n\left(\frac{1}{3}+x\right)_n\left(\frac{2}{3}+x\right)_n}\left(\frac{16}{27}\right)^n\frac{11(n+x)-3}{(n+x)^3}$$
$$=\frac{6(3x-1)(3x-2)}{x^3(2x-1)}\sum_{k=0}^{\infty}\frac{\left(\frac{1}{2}\right)_k\left(\frac{3}{2}-x\right)_k}{\left(\frac{1}{2}+x\right)_k^2}.$$

Taking $x=1$ we again obtain (14).

Acknowledgment

I thank W. Zudilin for encouraging me to use the Barnes integral representation to obtain the sum of "divergent" Ramanujan-type series. I thank G. Almkvist for suggesting to me to integrate on the left side to get identities with convergent series. I am also grateful to Jonathan Sondow for several helpful comments.

References

[1] G. Almkvist *Glaisher's formulas for* $1/\pi^2$ *and some generalizations*, (2011).

[2] W.N. Bailey, Generalized hypergeometric series, Cambridge Math. Tracts **32**, Cambridge Univ. Press, Cambridge, (1935); 2^{nd} reprinted edition, Stechert-Hafner, New York-London, (1964).

[3] D.H. Bailey, D. Borwein, J.M. Borwein, R.E. Crandall *Hypergeometric forms for Ising-class integrals*, Experiment. Math. **16**, 257-276, (2007).

[4] Ekhad, S.B., Zeilberger D.: *A WZ proof of Ramanujan's formula for* π. In Rassias, J.M. (ed.). Geometry, Analysis and Mechanics. World Scientific, Singapore, 107-108, (1994); also available at arXiv:math/9306213v1. (The coauthor EKHAD is a Maple package written by D. Zeilberger).

[5] J. Guillera, *Some binomial series obtained by the WZ-method.* Adv. in Appl. Math. **29**, 599-603, (2002); arXiv:math/0503345.

[6] J. Guillera, *Generators of Some Ramanujan Formulas*, Ramanujan J. **11**, 41-48, (2006).

[7] J. Guillera, *Hypergeometric identities for* 10 *extended Ramanujan-type series*, Ramanujan J., **15**, 219-234 (2008).

[8] J. Guillera, Series de Ramanujan: Generalizaciones y conjeturas. Ph.D. Thesis, University of Zaragoza, Spain, (2007).

[9] J. Guillera, *On WZ-pairs which prove Ramanujan series*, Ramanujan J, **22**, 249-259, (2008); arXiv:0904.0406.

[10] J. Guillera, *A new Ramanujan-like series for* $1/\pi^2$, Ramanujan J. **26**, 369-374, (2011); arXiv:1003.1915.

[11] J. Guillera, *More hypergeometric identities related to Ramanujan-type series*, To appear in Ramanujan J.; arXiv:1003.1915.

[12] J. Guillera and W. Zudilin, *"Divergent" Ramanujan-type supercongruences*, Proc. of the Amer. Math. Soc. **140**, 765-777, (2012); arXiv:1004.4337.

[13] M. Petkovšek, H. S. Wilf, D. Zeilberger, A=B, A K. Peters, Ltd., (1996); also available at http://www.math.upenn.edu/ wilf/AeqB.html.

[14] F. Stan, *On recurrences for Ising integrals*, Adv. in Appl. Math. **45**, 334-345, (2010).

[15] Z.W. Sun, *Supercongruences and Euler numbers*, Sci. China Math. **54**, 2509-2535, (2011); arXiv:1001.4453.

[16] E.T. Whittaker, G.N. Watson, A Course of Modern Analysis. Cambridge Univ. Press, (1927).

[17] H.S. Wilf, D. Zeilberger, *Rational functions certify combinatorial identities*, J. Amer. Math. Soc. **3**, 147-158, (1990) (winner of the Steele Prize).

E-mail address: `jguillera@gmail.com`

AV. CESÁREO ALIERTA, 31 ESC. IZDA 4^o–A, ZARAGOZA (SPAIN)